Thermodynamic Volume Product in Spherically Symmetric and Axisymmetric Spacetime

Parthapratim Pradhan*
Department of Physics, Vivekananda Satavarshiki Mahavidyalaya, West Midnapur 721513, India

arXiv:1611.04284v1 [gr-qc] 14 Nov 2016

Abstract

In this Letter, we compute particularly thermodynamic *volume product, volume sum, volume minus and volume division* for wide variety of spherically symmetric spacetime and axisymmetric spacetime in the frame work of *extended phase space*. We consider Einstein gravity as well as other than Einstein gravity i.e. *Hořava Lifshitz* gravity. We speculate that for spherically symmetric black holes the volume product is mass-independent both in Einstein gravity as well as Hořava Lifshitz gravity while the other combination is mass-dependent. For axisymmetric black hole spacetime in Einstein gravity all the combination is *mass-dependent*. There has been no chance to generate any combination of volume product is mass-independent. Interestingly, *only rotating BTZ black hole* in 3D provides the volume product formula is mass-independent i.e. *universal* and hence it is quantized.

1 Introduction

It has been examined by number of researchers that the area (or entropy) product of various spherically symmetric and axisymmetric black holes are mass-independent (universal) [1, 2, 3, 5, 6, 7, 8, 9]. For instance, Ansorg and Hennig [1] demostrated that for a stationary and axisymmetric Einstein-Maxwell gravity the area product formula satisfied universal relation as

$$\mathcal{A}_{OH}\mathcal{A}_{IH} \;=\; (8\pi J)^2 + (4\pi Q^2)^2 \; . \tag{1}$$

where $\mathcal{A}_{OH}$ and $\mathcal{A}_{IH}$ are area of outer horizon (OH) or event horizon (EH) and inner horizon (IH) or Cauchy horizon (CH). J and Q are angular momentum and charge of the black hole (BH) respectively.

On the other hand, Cvetič et al. [2] extended this work for higher dimensions and showed for multihorizon BHs the area product formula should be quantized by satisfied the following relation:

$$\mathcal{A}_{OH}\mathcal{A}_{IH} \;=\; (8\pi\ell_{pl}^2)^2 N,\; N \in \mathbb{N} \; . \tag{2}$$

where ℓ_{pl} is the Planck length. This relation indicated that the product relation is indeed universal in nature. This is a very fascinatic topic of research since 2009.

Aspects of BH thermodynamic properties have started by the seminal work of Hawking and Page [10], they first showed the existence of a certain type of phase transition between small and large BHs in case of Schwarzschild-AdS BH. This phase transition is now called famous Hawking-Page phase transition. For charged AdS BH, the thermodynamic properties started by the work of Chamblin et al. [11, 12], where the authors demonstrated the critical behavior of an another type of Vander der Waals liquid-gas phase transition. This has been brought into a new form by Kubizňák and Mann [13] by studying the thermodynamic properties i.e. $P-V$ criticality of Reissner-Nordström AdS BH in the extended phase space. They determined the BH equation of state and computed critical exponent by using mean field theory and also computed other thermodynamic features.

Motivated by the above mentioned work and our previous investigation [14] where we have considered the extended phase space framework for wide variety of spherically symmetric AdS spacetime,

*ppradhan77@gmail.com

here we intend to extend our study for wide variety of spacetime by giving several examples including *3D static BTZ BH, Hořava Lifshitz BH and spinning BTZ BH.* In this framework, the cosmological constant treated as thermodynamic pressure P and its conjugate variable as thermodynamic volume $\mathcal{V}$ [15, 16, 17, 13]. They are defined as

$$P = -\frac{\Lambda}{8\pi} = \frac{3}{8\pi\ell^2} . \tag{3}$$

and

$$\mathcal{V} = \left(\frac{\partial M}{\partial P}\right)_{S,Q,J} . \tag{4}$$

This thermodynamic volume of spherically symmetric BH for OH reads

$$\mathcal{V}_{OH} = \frac{4}{3}\pi r_{OH}^3 = \frac{\mathcal{A}_{OH} r_{OH}}{3} . \tag{5}$$

where r_{OH} is OH radius. Similarly, this volume for IH should be

$$\mathcal{V}_{IH} = \frac{4}{3}\pi r_{IH}^3 = \frac{\mathcal{A}_{IH} r_{IH}}{3} . \tag{6}$$

It should be noted that the thermodynamic volume of CH can be obtain by using the symmetric properties [14] of OH radius r_{OH} and IH radius r_{IH} i.e.

$$\mathcal{V}_{IH} = V_{OH}|_{r_{OH}\leftrightarrow r_{IH}} . \tag{7}$$

The another important point in the extended phase space is that the ADM mass should be treated as the total enthalpy of the thermodynamic system i.e. $M = H = U + P\mathcal{V}$. Where U is thermal energy of the system [15]. Therefore the first law of BH thermodynamics in this phase space for any spherically symmetric spacetime for OH should be

$$dH = T_{OH} dS_{OH} + \mathcal{V}_{OH} dP + \Phi_{OH} dQ . \tag{8}$$

where the quantities T_{OH}, S_{OH} and Φ_{OH} are the BH temperature, entropy and electric potential of OH, and Q is charge of the BH.

Analogously, the first law of BH mechanics for IH must be

$$dH = T_{IH} dS_{IH} + \mathcal{V}_{IH} dP + \Phi_{IH} dQ . \tag{9}$$

where the quantities T_{IH}, S_{IH} and Φ_{IH} are the corresponding BH temperature, entropy and electric potential defined on the IH.

When we add the rotation the first law of BH thermodynamics in the extended phase space for axisymmetric spacetime for OH becomes

$$dH = T_{OH} dS_{OH} + \mathcal{V}_{OH} dP + \Phi_{OH} dQ + \Omega_{OH} dJ . \tag{10}$$

where Ω_{OH} and J are the angular velocity defined on OH and the angular momentum of BH. For IH, the first law becomes

$$dH = T_{IH} dS_{IH} + \mathcal{V}_{IH} dP + \Phi_{IH} dQ + \Omega_{IH} dJ . \tag{11}$$

where Ω_{IH} is the angular velocity defined on IH.

However in this work, we intend to extend our study by computing the volume product, volume sum, volume minus and volume division in the *extended phase space* for various spherically symmetric BHs and axisymmetric BHs. By evaluating these quantities we prove that for spherically symmetric

BHs [1] the thermodynamic volume products are mass-independent therefore they behaves as universal character by its own features. While the axisymmetric BHs showing *no* universal behavior. Interestingly, only rotating BTZ BH shows the mass-independent features. Thus *only* axisymmetric BHs in 3D provided the universal character of thermodynamic volume product.

In our previous investigation [8, 9], we computed the BH area (or entropy) product, BH temperature product, Komar energy product and specific heat product for various BHs. Besides the area (or entropy) product, it should be important to study whether the thermodynamic *volume product, volume sum, volume minus and volume division* for all the horizons are universal or not. This is main motivation behind this work. Whether they should be quantized or not.

The structure of the paper is as follows. In Sec. 2, we shall compute various thermodynamic volume products for spherically symmetric BHs and conclude that the product is mass-independent. In Sec. 3, we compute various thermodynamic volume products for axisymmetric spacetime and conclude that the product is mass-dependent. Interestingly, for the spinning BTZ BH the said volume product is *mass-independent*.

2 Spherically Symmetric BH

In this section we should considered only the various spherically symmetric BHs.

2.1 Reissner Nordström BH:

We begin with charged BH with zero cosmological constant which is a solution of Einstein equation. The OH radius and IH radius reads

$$r_{OH} = M + \sqrt{M^2 - Q^2} \tag{12}$$

$$r_{IH} = M - \sqrt{M^2 - Q^2}\ . \tag{13}$$

where M and Q denotes the mass and charge of BH respectively. When $M^2 > Q^2$, it descibes a BH otherwise it has a naked singularity. The thermodynamic volume for OH and IH should read

$$\mathcal{V}_{OH} = \frac{4}{3}\pi r_{OH}^3 \tag{14}$$

$$\mathcal{V}_{IH} = \frac{4}{3}\pi r_{IH}^3\ . \tag{15}$$

The thermodynamic volume[2] product for OH and IH should be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{16}{9}\pi^2 Q^6\ . \tag{16}$$

It is indeed mass-independent therefore it is universal in character and also quantized.

The volume sum for OH and IH is calculated to be

$$\mathcal{V}_{OH} + \mathcal{V}_{IH} = \frac{32}{3}\pi M^3\left(1 - \frac{3}{4}\frac{Q^2}{M^2}\right)\ . \tag{17}$$

Similarly, one can compute the volume minus for OH and IH as

$$\mathcal{V}_{OH} - \mathcal{V}_{IH} = \frac{32}{3}\pi M^2\sqrt{M^2 - Q^2}\left(1 - \frac{Q^2}{4M^2}\right)\ . \tag{18}$$

[1] except isolated Schwarzschild BH, Schwarzschild-AdS BH and static BTZ BH

[2] In the limit $Q = 0$, one obtains the thermodynamic volume for Schwarzschild BH. Since in this case the BH has only OH located at $r_{OH} = 2M$. Therefore the volume should be $\mathcal{V}_{OH} = \frac{32}{3}\pi M^3$. Thus for an isolated Schwarzschild BH the thermodynamic volume should be mass dependent therefore it is not universal and not quatized in nature by its own character.

and the volume division should be

$$\frac{\mathcal{V}_{OH}}{\mathcal{V}_{IH}} = \left(\frac{M+\sqrt{M^2-Q^2}}{M-\sqrt{M^2-Q^2}}\right)^3 . \tag{19}$$

It follows from the calculation that all these quantities are mass dependent so they are not universal in nature by its own right. From Eq. (18) and Eq. (19), we can easily see that in the extremal limit $M^2 = Q^2$, one obtains $\mathcal{V}_{OH} = \mathcal{V}_{IH}$. This is a new condition of extreme limit in spherically symmetric cases.

2.2 Hořava Lifshitz BH:

Now we consider another spherically symmetric BH in Hořava Lifshitz (HL) [24, 25, 26] gravity other than Einstein gravity but it reduces to Einstein's gravity at large scales for dynamical coupling constant $\lambda = 1$. It is a UV complete non-relativistic renormalizable theory of gravity. One possible class of solution in HL gravity is Kehagias-Sfetsos (KS) BH [23]. The OH radius and IH radius should read

$$r_{OH} = M + \sqrt{M^2 - \frac{1}{2\omega}} \tag{20}$$

$$r_{IH} = M - \sqrt{M^2 - \frac{1}{2\omega}} . \tag{21}$$

where M and ω denotes the mass and coupling constant of BH respectively. When $M^2 > \frac{1}{2\omega}$, it descibes a BH and when $M^2 < \frac{1}{2\omega}$, it descibes a naked singularity.

The thermodynamic volume product for KS BH should be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{2\pi^2}{9\omega^3} . \tag{22}$$

It indicates that it is mass-independent therefore it is universal in nature and it also be quantized. We do not calculate other possible combination because it is clear that these combinations are surely mass dependent as we have seen in case of RN BH.

2.3 Non-rotating BTZ BH:

It is interesting to study another special class of BH in 3D gravity is static BTZ BH. The BH OH is at $r_{OH} = \sqrt{8G_3M\ell}$. G_3 is 3D Newtonian constant. Interestingly, the thermodynamic volume for 3D static BTZ BH computed in [17]

$$\mathcal{V}_{OH} = \pi r_{OH}^2 = 8\pi G_3 M \ell^2 \tag{23}$$

This is an isolated case and the thermodynamic volume is mass dependent thus it is not quantized and not universal. where $\Lambda = -\frac{1}{\ell^2}$ is cosmological constant.

2.4 Schwarzschild-AdS BH:

This BH is a solution of Einstein equation. The BH has a single physical horizon [6] is at

$$r_{OH} = \frac{2\ell}{\sqrt{3}} sinh\left[\frac{1}{3} sinh^{-1}\left(3\sqrt{3}\frac{M}{\ell}\right)\right] \tag{24}$$

where $\Lambda = -\frac{3}{\ell^2}$ is cosmological constant. The thermodynamic volume is computed to be

$$\mathcal{V}_{OH} = \frac{4}{3}\pi r_{OH}^3 = \frac{4}{3}\pi\left(\frac{2\ell}{\sqrt{3}} sinh\left[\frac{1}{3} sinh^{-1}\left(3\sqrt{3}\frac{M}{\ell}\right)\right]\right)^3 . \tag{25}$$

Since it is isolated case and the thermodynamic volume is mass dependent therefore it is not universal nor does it quantized. We do not considered the other AdS spacetime because it has already been discussed in [14]. Now we are moving to axisymmetric spacetime. To see what happens there?

3 Axisymmetric Spacetime

In this section we have considered only the various axisymmetric BHs. It is easy to compute volume products for spherically symmetric cases bacause of $\mathcal{V}_{OH} \propto \mathcal{A}_{OH} r_{OH}$ for OH and $\mathcal{V}_{IH} \propto \mathcal{A}_{IH} r_{IH}$ for IH. While for axisymmetric spacetime this proportionality is quite different because here the spin parameter is present. Now see what happens in this case by starting with Kerr BH.

3.1 Kerr BH:

The Kerr BH is a solution of Einstein equation. The OH radius and IH radius reads for this BH should read

$$r_{OH} = M + \sqrt{M^2 - a^2} \tag{26}$$

$$r_{IH} = M - \sqrt{M^2 - a^2}\ . \tag{27}$$

where $a = \frac{J}{M}$. J is angular momentum of the BH. When $M^2 > a^2$, it descibes a BH while $M^2 < a^2$ it descibes a naked singularity. The thermodynamic volume for OH [19] and IH [14] becomes

$$\mathcal{V}_{OH} = \frac{\mathcal{A}_{OH} r_{OH}}{3}\left[1 + \frac{a^2}{2r_{OH}^2}\right] \tag{28}$$

$$\mathcal{V}_{IH} = \frac{\mathcal{A}_{IH} r_{IH}}{3}\left[1 + \frac{a^2}{2r_{IH}^2}\right]\ . \tag{29}$$

The thermodynamic volume product of Kerr BH for OH and IH is calculated to be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{128}{9}\pi^2 J^2 M^2 \left(1 + \frac{a^2}{8M^2}\right)\ . \tag{30}$$

The volume sum for OH and IH is

$$\mathcal{V}_{OH} + \mathcal{V}_{IH} = \frac{32}{3}\pi M^3 \left(1 - \frac{a^2}{4M^2}\right)\ . \tag{31}$$

Similarly, the volume minus for OH and IH is

$$\mathcal{V}_{OH} - \mathcal{V}_{IH} = \frac{32}{3}\pi M^2 \sqrt{M^2 - a^2}\ . \tag{32}$$

and the volume division is

$$\frac{\mathcal{V}_{OH}}{\mathcal{V}_{IH}} = \left(\frac{4M^2 - a^2 + 4M\sqrt{M^2 - a^2}}{4M^2 - a^2 - 4M\sqrt{M^2 - a^2}}\right)\ . \tag{33}$$

It indicates that the volume product, volume sum, volume minus and volume division for Kerr BH is mass dependent. Therefore the product, the sum, the minus and the division all are *not* universal.

3.2 Kerr-AdS BH:

The horizon function for Kerr-AdS BH [22] is given by

$$\Delta_r = \left(r^2 + a^2\right)\left(1 + \frac{r^2}{\ell^2}\right) - 2Mr = 0\ . \tag{34}$$

which gives the quartic order ofhorizon equation

$$\frac{r^4}{\ell^2} + \left(1 + \frac{a^2}{\ell^2}\right) r^2 - 2mr + a^2 = 0\ . \tag{35}$$

The quantities m and a are related to the parameters mass M and angular momentum J as follows

$$m = M\Xi^2,\ a = \frac{J}{m}\Xi^2 \tag{36}$$

where $\Xi = 1 - \frac{a^2}{\ell^2}$. To obtain the roots of Eq. (35) we apply the Vieta's theorem, we find

$$\sum_{i=1}^{4} r_i = 0\ . \tag{37}$$

$$\sum_{1\leq i<j\leq 4} r_i r_j = \ell^2\left(1+\frac{a^2}{\ell^2}\right)\ . \tag{38}$$

$$\sum_{1\leq i<j<k\leq 4} r_i r_j r_k = 2m\ell^2\ . \tag{39}$$

$$\prod_{i=1}^{4} r_i = a^2\ell^2\ . \tag{40}$$

There is at least two real zeros of the Eq. (35) which is OH radius and IH radius. After some algebraic computation we have

$$r_{OH} + r_{IH} = \frac{2m\ell^2}{a^2+\ell^2+r_{OH}^2+r_{IH}^2}, \tag{41}$$

$$r_{OH}r_{IH} = \frac{a^2\ell^2-(r_{OH}r_{IH})^2}{a^2+\ell^2+r_{OH}^2+r_{IH}^2}\ . \tag{42}$$

The area of the BH for OH is

$$\mathcal{A}_{OH} = \frac{4\pi\left(r_{OH}^2+a^2\right)}{\Xi}\ . \tag{43}$$

and for IH is

$$\mathcal{A}_{IH} = \frac{4\pi\left(r_{IH}^2+a^2\right)}{\Xi}\ . \tag{44}$$

The thermodynamic volume for OH [17, 18] becomes

$$\mathcal{V}_{OH} = \frac{2\pi\left[(r_{OH}^2+a^2)(2r_{OH}^2\ell^2+a^2\ell^2-r_{OH}^2a^2)\right]}{3r_{OH}\ell^2\Xi^2}\ . \tag{45}$$

and we derive the thermodynamic volume for IH becomes

$$\mathcal{V}_{IH} = \frac{2\pi\left[(r_{IH}^2+a^2)(2r_{IH}^2\ell^2+a^2\ell^2-r_{IH}^2a^2)\right]}{3r_{IH}\ell^2\Xi^2}\ . \tag{46}$$

The thermodynamic volume product for Kerr-AdS BH is calculated to be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{4\pi^2\{r_{OH}^2r_{IH}^2+a^2\left(r_{OH}^2+r_{IH}^2\right)+a^4\}}{9\Xi^4 r_{OH}r_{IH}} \times \left[3r_{OH}^2r_{IH}^2+2a^2\left(r_{OH}^2+r_{IH}^2\right)+\Xi^2r_{OH}^2r_{IH}^2\right]\ . \tag{47}$$

Using Eq. (41), Eq. 42, Eq. (43) and Eq. (44), we observe that there is no way to eliminate the mass parameter from Eq. (47) therefore the volume product for Kerr-AdS BH is not mass-independent thus it is not universal and not quantized .

3.3 Kerr-Newman BH:

It is an axisymmetric solution of Einstein-Maxwell equations. The OH radius and IH radius for this BH becomes

$$r_{OH} = M + \sqrt{M^2 - a^2 - Q^2} \tag{48}$$
$$r_{IH} = M - \sqrt{M^2 - a^2 - Q^2}\ . \tag{49}$$

The thermodynamic volume for OH [17] is

$$\mathcal{V}_{OH} \;=\; \frac{2\pi\left[(r_{OH}^2 + a^2)(2r_{OH}^2 + a^2) + a^2Q^2\right]}{3r_{OH}}\ . \tag{50}$$

and we derive that the thermodynamic volume for IH is

$$\mathcal{V}_{IH} \;=\; \frac{2\pi\left[(r_{IH}^2 + a^2)(2r_{IH}^2 + a^2) + a^2Q^2\right]}{3r_{IH}}\ . \tag{51}$$

The thermodynamic volume product for KN BH is computed to be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{16\pi^2}{9(a^2+Q^2)}\left[J^2(8J^2 + a^4 - a^2Q^2 - 2Q^4 + 8M^2Q^2) + Q^4(a^2+Q^2)^2\right]\ . \tag{52}$$

It also indicates that the thermodynamic volume for KN BH is mass dependent. Thus the volume product is not universal for any axisymmetric spacetime. In the appropriate limit i.e. when $a = J = 0$, one obtains the thermodynamic volume product for Reissner Nordström BH and when $Q = 0$, one obtains the volume product for Kerr BH.

3.4 Kerr-Newman-AdS BH:

The horizon function for Kerr-Newman-AdS BH [21] reads

$$\Delta_r \;=\; \left(r^2 + a^2\right)\left(1 + \frac{r^2}{\ell^2}\right) - 2Mr + q^2 = 0\ . \tag{53}$$

which has the quartic order of horizon equation

$$\frac{r^4}{\ell^2} + \left(1 + \frac{a^2}{\ell^2}\right)r^2 - 2mr + a^2 + q^2 \;=\; 0\ . \tag{54}$$

The quantity q is related to the charge parameter Q as

$$q \;=\; Q\Xi \tag{55}$$

To determine the roots of Eq. (54) again we apply the Vieta's rule then one obtains

$$\sum_{i=1}^{4} r_i \;=\; 0\ . \tag{56}$$
$$\sum_{1\le i<j\le 4} r_i r_j \;=\; \ell^2\left(1 + \frac{a^2}{\ell^2}\right)\ . \tag{57}$$
$$\sum_{1\le i<j<k\le 4} r_i r_j r_k \;=\; 2m\ell^2\ . \tag{58}$$
$$\prod_{i=1}^{4} r_i \;=\; (a^2 + q^2)\ell^2\ . \tag{59}$$

Similarly, there is at least two real zeros of the Eq. (54) which is OH radius and IH radius. After some algebraic derivation one get

$$r_{OH}+r_{IH} = \frac{2m\ell^2}{a^2+\ell^2+r_{OH}^2+r_{IH}^2}, \tag{60}$$

$$r_{OH}r_{IH} = \frac{(a^2+q^2)\ell^2-(r_{OH}r_{IH})^2}{a^2+\ell^2+r_{OH}^2+r_{IH}^2}\ . \tag{61}$$

The area of this BH for OH is

$$\mathcal{A}_{OH} = \frac{4\pi\left(r_{OH}^2+a^2\right)}{\Xi}\ . \tag{62}$$

and for IH is

$$\mathcal{A}_{IH} = \frac{4\pi\left(r_{IH}^2+a^2\right)}{\Xi}\ . \tag{63}$$

The thermodynamic volume for OH [17, 18] becomes

$$\mathcal{V}_{OH} = \frac{2\pi\left[(r_{OH}^2+a^2)(2r_{OH}^2\ell^2+a^2\ell^2-r_{OH}^2a^2)+\ell^2q^2a^2\right]}{3r_{OH}\ell^2\Xi^2}\ .$$

and we derive the thermodynamic volume for IH becomes

$$\mathcal{V}_{IH} = \frac{2\pi\left[(r_{IH}^2+a^2)(2r_{IH}^2\ell^2+a^2\ell^2-r_{IH}^2a^2)+\ell^2q^2a^2\right]}{3r_{IH}\ell^2\Xi^2}\ .$$

The thermodynamic volume product for Kerr-Newman-AdS BH is computed to be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{4\pi^2}{9\Xi^4r_{OH}r_{IH}}\times$$

$$\left[\{r_{OH}^2r_{IH}^2+a^2\left(r_{OH}^2+r_{IH}^2\right)+a^4\}^2+\{r_{OH}^2r_{IH}^2+a^2\left(r_{OH}^2+r_{IH}^2\right)+a^4\}\{2r_{OH}^2r_{IH}^2+a^2\left(r_{OH}^2+r_{IH}^2\right)\}\right]$$

$$+\frac{4\pi^2\left[a^2q^2\{r_{OH}^4+r_{IH}^4+2a^2\left(r_{OH}^2+r_{IH}^2\right)+2a^4\}+\Xi^2r_{OH}^2r_{IH}^2\{r_{OH}^2r_{IH}^2+a^2\left(r_{OH}^2+r_{IH}^2\right)+a^4\}\right]}{9\Xi^4r_{OH}r_{IH}}$$

$$+\frac{4\pi^2\left[a^2q^2\Xi\{r_{OH}^4+r_{IH}^4+a^2\left(r_{OH}^2+r_{IH}^2\right)\}+a^4q^4\right]}{9\Xi^4r_{OH}r_{IH}}\ . \tag{64}$$

Again using Eq. (60), Eq. 61, Eq. (62) and Eq. (63), we speculate that there has been no chance to eliminate the mass parameter from Eq. (64) thus the volume product for Kerr-Newman-AdS BH is not mass-independent therefore it is not universal and not quantized .

3.5 Spinning BTZ BH:

The metric for rotating BTZ BH [27] in $2+1$ dimension is given by

$$ds^2 = -\left(\frac{r^2}{\ell^2}+\frac{J^2}{4r^2}-M\right)dt^2+\frac{dr^2}{\left(\frac{r^2}{\ell^2}+\frac{J^2}{4r^2}-M\right)}$$

$$+r^2\left(-\frac{J}{2r^2}dt+d\phi\right)^2\ . \tag{65}$$

where M and J represents the ADM mass, and the angular momentum of the BH. $-\Lambda=\frac{1}{\ell^2}=8\pi PG_3$ denotes the cosmological constant. Here we have set $8G_3=1=c=\hbar=k$. When $J=0$, one obtains the static BTZ BH.

The BH OH radius and IH radius are [27, 20]:

$$r_{OH} = \sqrt{\frac{M\ell^2}{2}\left(1+\sqrt{1-\frac{J^2}{M^2\ell^2}}\right)}. \tag{66}$$

$$r_{IH} = \sqrt{\frac{M\ell^2}{2}\left(1-\sqrt{1-\frac{J^2}{M^2\ell^2}}\right)}. \tag{67}$$

The thermodynamic volume for 3D spinning BTZ BH for OH and IH:

$$\mathcal{V}_{OH} = \left(\frac{\partial M}{\partial P}\right)_J = \pi r_{OH}^2 \tag{68}$$

$$\mathcal{V}_{IH} = \left(\frac{\partial M}{\partial P}\right)_J = \pi r_{IH}^2 \tag{69}$$

The thermodynamic volume product is computed to be

$$\mathcal{V}_{OH}\mathcal{V}_{IH} = \frac{\pi^2 J^2 \ell^2}{4}\,. \tag{70}$$

Interestingly, the thermodynamic volume product for rotating BTZ BH is *mass-independent i.e. universal* and it is also quantized. This is the only example for *rotating* cases, the volume product is universal. This is an interesting result of this work.

4 Discussion:

In this work, we demonstrated the thermodynamic products particularly thermodynamic volume products of spherically symmetric spacetime and axisymmetric spacetime by incorporating the extended phase-space formalism. In this formalism, the cosmological constant should be considered as a thermodynamic pressure and its conjugate parameter as thermodynamic volume. Additionaly, the mass parameter should be treated as enthalpy of the system rather than internal energy. Then in this phase space the first law of BH thermodynamics should be satisfied for both OH and IH.

We explicily computed the thermodynamic volume products both for OH and IH of several classes of spherically symmetric and axisymmetric BHs. Such products are *mass independent* for spherically symmetric cases (except isolated Schwarzschild BH, Schwarzschild-AdS BH and static BTZ BH) because of $\mathcal{V}_{OH} \propto \mathcal{A}_{OH} r_{OH}$ for OH and $\mathcal{V}_{IH} \propto \mathcal{A}_{IH} r_{IH}$ for IH. Axisymmetric spacetime does not satisfied this proportionality beacuse of introduction of the spin parameter thus they showed *no* mass-independent features except the rotating BTZ BH, the only axisymmetric spacetime in $3D$ showed *universal* features thus it has been quantized in this sense. We also computed thermodynamic volume sum but they are always mass dependent so they are not universal as well as they are not quantized. Besides area (or entropy) products the thermodynamic volume products of OH and IH that we have investigated in this work in some cases (but not all) further provide some universal properties of the BH which gives some insight of microscopic origin of BH entropy both outer and inner.

References

[1] M. Ansorg and J. Hennig, *Phys. Rev. Lett.* **102**, 221102 (2009).

[2] M. Cvetič et al., *Phys. Rev. Lett.* **106**, 121301 (2011).

[3] A. Castro and M. J. Rodriguez, *Phys. Rev. D* **86**, 024008 (2012).

[4] A. Castro et al. , *JHEP* **07**, 164 (2013).

[5] S. Detournay, *Phys. Rev. Lett.* **109**, 031101 (2012).

[6] M. Visser, *Phys. Rev. D* **88**, 044014 (2013).

[7] J. Hennig, *Class. Quant. Grav.* **31** 135005 (2014).

[8] P. Pradhan, *Eur. Phys. J.* **C 74**, 2887 (2014).

[9] P. Pradhan, *Phys. Lett. B* **747**, 64 (2015).

[10] S. Hawking and D. Page, *Commun. Math. Phys.* **87**, 577 (1983).

[11] A. Chamblin et al., *Phys.Rev. D 60*, 064018 (1999).

[12] A. Chamblin et al., *Phys. Rev. D 60*, 104026 (1999).

[13] D. Kubizňák and R. B. Mann, *JHEP* **1207**, 033 (2012).

[14] P. Pradhan, *Int. J. Mod. Phys. D* **26**, 1750010 (2017).

[15] D. Kastor et al., *Class. Quant. Grav.* **26**, 195011 (2009).

[16] B. P. Dolan, *Class. Quant. Grav.* **28**, 125020 (2011) .

[17] B. P. Dolan, *Class. Quant. Grav.* **28**, 235017 (2011).

[18] M. Cvetič et al., *Phys. Rev.* **D84**, 024037 (2011).

[19] N. Altamirano et al. , *Galaxies* **2**, 89 (2014).

[20] P. Pradhan, *JETP Letters.* **102**, 427 (2015).

[21] M. M. Caldrelli et al., *Class. Quant. Grav.* **17**, 399 (2000).

[22] B. Carter, *Commun. Math. Phys.* **10**, 280 (1968).

[23] A. Kehagias and K. Sfetsos, *Phys. Lett.* B 678, 123 (2009).

[24] P. Horava, *Phys. Rev. Lett.* **102**, 161301 (2009).

[25] P. Horava, *Phys. Rev.* **D 79**, 084008 (2009).

[26] P. Horava, *JHEP* **0903**, 020 (2009).

[27] M. Bañados et al., *Phys. Rev. Lett.* **69** 1849 (1992).